

DOSE PREPARATION WORKSHEET

Bebtelovimab 175mg IV Push Injection



INTRAVENOUS PUSH PREPARATION

Gather supplies:

- 1 vial of **Bebtelovimab 175mg/2 mL**
- 3 mL syringe
- 0.9% Sodium Chloride flush
- 18 or 20 gauge needles (1)
- Alcohol pads

Remove 1 vial of **Bebtelovimab 175mg/2 mL** from the refrigerator. Allow to equilibrate to room temperature for approximately 20 minutes before preparation. Do not expose to direct heat. Do not shake vial.

Don PPE: mask, gloves, gown and eye protection.

Inspect the **Bebtelovimab 175mg/2 mL** vial visually for particulate matter and discoloration. Bebtelovimab is clear to opalescent and colorless to slightly yellow to slightly brown solution. Discard the vial if the solution is cloudy, discolored or visible particles are observed.

Withdraw 2 mL from the vial of **Bebtelovimab 175mg/2 mL** into a 3mL syringe.
Dual nurse verification required to validate dose.

Discard any product remaining in the vial.

Adhere pharmacy label to syringe: Bebtelovimab 175mg/2 mL

ADMINISTRATION

Verify patient using 2 identifiers

Obtain vital signs: blood pressure, pulse, O2 saturation and temperature.

- ✓ Record prior to infusion; at 15 minutes; 60 minutes after injection before patient is discharged

Discard needle and attach syringe to the IV hub for injection

Administer the entire contents of the syringe via *IV Push injection* over at least *30 seconds*.

Flush the IV hub with 10 mL 0.9% Sodium Chloride to ensure full dose delivered.

Monitor patient post injection for 1 hour, obtain vital signs 60 minutes after the injection.

Document patient's tolerance to bebtelovimab in progress note.

Instruct patient to report any reactions experienced after discharge to PCP. PCP to complete FDA MedWatch form should this occur.

Pharmacy Label

Patient Name: _____ DOB: _____ Bebtelovimab 175mg/ 2mL solution <i>Inject 175mg IV Push over at least 30 seconds</i> Lot number _____ Manuf Exp _____ Prepared Date/Time _____ Exp Time _____ Initials _____
--